

Nova Science Publishers, Inc.
African Diaspora Journal of Mathematics
Volume 00, No. 00, p. 000–000 (2004)
ISSN 1539-854X Copyright ©2004
www.african-j-math.org

Non abelian cohomology: the point of view of gerbed tower

Aristide Tsemo*

February 1, 2008

Abstract

We define in this paper the notion of gerbed tower. This enables us to interpret geometrically cohomology classes without using the notion of n -category. We use this theory to study sequences of affine maps between affine manifolds, and the cohomology of manifolds.

keywords gerbes, non abelian cohomology.

Classification A.M.S. 18D05, 57R20.

1 Introduction

In mathematics, a theory is defined by axioms which describe relations between elements of a set. The purpose of mathematicians is to classify these elements by defining structures modelled on reference objects. In geometry, a structure modelled on the space L , is defined on a topological set N , by a Čech 0-chain whose boundary reflects properties of L . For example, a n -differentiable manifold is defined by an atlas $(U_i)_{i \in I}$, and charts $\phi_i : U_i \rightarrow \mathbb{R}^n$, such that $\phi_j \circ \phi_i^{-1}$ is a differentiable map, here the model L is $\mathbb{R}^n$, and the property reflected is the differentiability. The manifold is obtained by gluing the sets $\phi_i(U_i)$ using the 1-cocycle $h_{ij} = \phi_j \circ \phi_i^{-1}$. Alternatively, a structure is defined by gluing a family of sets N_i using a cocycle h_{ij} . Often the sets N_i are related to the model in the sense that each of them is endowed with a L -structure. The natural problem to determine if a given topological space N can be endowed with a structure modelled on L , leads to the notion of sheaf of categories. When the structure exists locally, that is when there exists an open cover $(N_i)_{i \in I}$ of N , such that each N_i is endowed with a L -structure, the existence of the L -structure on N is equivalent to determine whether the cohomology class of a 2-Čech cocycle is trivial. This has motivated the definition of a 2-structure called gerbe, which is classified in geometry by a 2-Čech cocycle. The natural problem which occurs is to provide geometric conditions which insure a 2-Čech chain to define a 2-type structure, ..., a n -Čech chain to define a n -type structure. On this purpose, one needs to give a geometric interpretation of Čech cohomology classes. Unfortunately, the notion of n -category needed to define n -structures is not well-understood. The main goal of this paper is to interpret Čech classes geometrically, by defining the notion of commutative n -gerbed tower. These are sequences of 2-categories $F_n \rightarrow F_{n-1} \dots F_2 \rightarrow F_1$, where F_1 is a gerbe

*Department, of Mathematics, St George University of Toronto Toronto, ON Canada M5S 3G3. (tsemo58@yahoo.ca).

defined on a topos N . A commutative n -gerbed tower satisfy conditions which allow to attach to it a family of p -Čech cohomology classes $([f_2], \dots, [f_{n+1}])$, where $[f_p] \in H^p(N, L_{p-1})$, and L_p are commutative sheaves defined on N . This notion represents geometrically the connecting morphism in cohomology. More precisely we have:

Theorem 4.2.6.

Let $u_n = F_n \rightarrow F_{n-1} \dots F_2 \rightarrow F_1$ be a commutative n -gerbed tower defined on a topos to which is associated the family of cohomology classes $([f_2], \dots, [f_{n+1}])$. Suppose that there exists an exact sequence of sheaves $0 \rightarrow L_{n+1} \rightarrow L'_{n+1} \rightarrow L_n \rightarrow 0$, then the family of cohomology classes $([f_2], \dots, [f_{n+2}])$ where $[f_{n+2}]$ is the image of $[f_{n+1}]$ by the connecting morphism $H^{n+1}(N, L_n) \rightarrow H^{n+2}(N, L_{n+1})$, is associated to a $n+1$ -gerbed tower.

An example of a n -gerbed tower appears in the theory of affine manifolds. An affine manifold (N, ∇_N) is a differentiable manifold N , endowed with a connection ∇_N , whose curvature and torsion forms vanish identically. We say that the n -dimensional affine manifold (N, ∇_N) , is complete, if and only if it is the quotient of the affine space $\mathbb{R}^n$, by a subgroup Γ_N of $Aff(\mathbb{R}^n)$ which acts properly and freely on $\mathbb{R}^n$. L. Auslander has conjectured that the fundamental group of a compact and complete affine manifold is polycyclic. In [26] we have conjectured that a finite Galois cover of a compact and complete affine manifold is the domain of a non trivial affine map. This leads to the following problem: classify sequences $(N_n, \nabla_{N_n}) \rightarrow \dots \rightarrow (N_1, \nabla_{N_1})$ where each map $f_i : (N_{i+1}, \nabla_{N_{i+1}}) \rightarrow (N_i, \nabla_{N_i})$ is an affine fibration whose domain is compact and complete: This means that f_i is a surjective map and each affine manifold (N_i, ∇_{N_i}) is a compact and complete affine manifold. The classification of affine fibrations has been done using gerbe theory (see [28]). It is normal to think that composition sequences of affine manifolds are related to n -gerbes. We define a n -gerbed tower which appears naturally in this context.

Characteristic classes are used in mathematics to study many objects, for example, Witten has used characteristic classes to study the Jones polynomial. This shows the necessity to give a geometric interpretation of characteristic classes. On this purpose, we have to interpret geometrically the integral cohomology of a differentiable manifold N . The theory of Kostant-Weil gives an interpretation of the group $H^2(N, \mathbb{Z})$: It is the set of equivalence classes of complex line bundles over N . In [5], is given an interpretation of $H^3(N, \mathbb{Z})$ in terms of equivalence classes of Dixmier-Douady groupoids. Our theory enables us to interpret a subgroup of $H^{n+2}(N, \mathbb{Z})$ as the set of equivalence classes of a family of n -gerbed towers.

This is the plan of our paper:

1. Introduction.
2. The notion of gerbe.
3. Notations
4. The notion of gerbed towers.
5. Spectral sequences and gerbed towers.
6. Applications of gerbed towers to affine geometry.
7. Interpretation of the integral cohomology of a manifold.
- 8 A definition of a notion of sheaf of n -categories.

2 The notion of gerbe.

In this part we present the notion of gerbe studied by Giraud [11].

Definitions 2.0.1.

Let E be a category, a **sieve** is a subclass R of the class of objects $Ob(E)$ of E such that if $f : X \rightarrow Y$ is a map of E , such that $Y \in R$, then $X \in R$.

Let $f : E' \rightarrow E$ be a functor, and R a sieve of E , we denote by R^f , the sieve defined by $R^f = \{X \in Ob(E') : f(X) \in R\}$.

For each object T of E , we denote by E_T , the category whose objects are arrows $u : U \rightarrow T$, a morphism of E_T between $u_1 : U_1 \rightarrow T$, and $u_2 : U_2 \rightarrow T$, is a map $h : U_1 \rightarrow U_2$ such that $u_2 \circ h = u_1$ •

Definition 2.0.2.

A **topology** on E is defined as follows: to each object T of E , we associate a non empty set $J(T)$ of sieves of the category E_T of E , above T such that:

- (i) For each map $f : T_1 \rightarrow T_2$, and for each element R of $J(T_2)$, $R^f \in J(T_1)$. (The morphism f induces a functor between E_{T_1} and E_{T_2} abusively denoted f).
- (ii) The sieve R of E_T is an element of $J(T)$, if for every map $f : T' \rightarrow T$ of E , $R^f \in J(T')$ •

A category endowed with a topology is called a site.

Definitions 2.0.3.

A **sheaf of sets** L defined on the category E endowed with the topology J , is a contravariant functor $L : E \rightarrow Set$, where Set is the category of sets, such that for each object U of E , and each element R of $J(U)$, the natural map:

$$L(U) \longrightarrow \lim(L \mid R)$$

is bijective, where $(L \mid R)$ is the correspondence defined on R by $(L \mid R)(f) = L(T)$ for each map $f : T \rightarrow U$ in R .

Let $h : F \rightarrow E$ be a functor, for each object U of E , we denote by F_U the subcategory of F defined as follows: an object T of F_U is an object T of F such that $h(T) = U$. A map $f : T \rightarrow T'$ between a pair of objects T and T' of F_U , is a map of F such that $h(f)$ is the identity of U . The category F_U is called **the fiber** of U . For each objects X , and Y of F_U , we will denote by $Hom_U(X, Y)$ the set of morphisms of F_U between X and Y •

Definitions 2.0.4.

Let $h : F \rightarrow E$ be a functor, $m : x \rightarrow y$ a map of F , and $f = h(m) : T \rightarrow U$ its projection by h . We will say that m is **cartesian**, or that m is the **inverse image** of f by h , or x is an inverse image of y by h , if for each element z of F_T , the map

$$Hom_T(z, x) \rightarrow Hom_f(z, y)$$

$$n \rightarrow mn$$

is bijective, where $Hom_f(z, y)$ is the set of maps $g : z \rightarrow y$ such that $h(g) = f$.

A functor $h : F \rightarrow E$ is a **fibred category** if and only if each map $f : T \rightarrow U$, has an inverse image, and the composition of two cartesian maps is a cartesian map.

We will say that the category is fibred in groupoids, if for each diagram

$$x \xrightarrow{f} z \xleftarrow{g} y$$

of F above the diagram of E ,

$$U \xrightarrow{\phi} W \xleftarrow{\psi} V$$

and for each map $m : U \rightarrow V$ such that $\psi m = \phi$, there exists a unique map $p : x \rightarrow y$, such that $gp = f$, and $h(p) = m$.

This implies that the inverse image is unique up to isomorphism.

Consider a map $\phi : U \rightarrow V$ of E , we can define a functor $\phi^* : F_V \rightarrow F_U$, such that for each object y of F_V , $\phi^*(y)$ is defined as follows: we consider a cartesian map $f : x \rightarrow y$ above ϕ and set $\phi^*(y) = x$. Remark that although the definition of $\phi^*(y)$ depends of the chosen inverse image f , the functors $(\phi\psi)^*$ and $\psi^*\phi^*$ are isomorphic •

Definitions 2.0.5.

A **section** of a fibred category $h : F \rightarrow E$, is a correspondence defined on the class of arrows of E as follows: to each map $f : U \rightarrow T$, we define a cartesian map: $l^f : x_U \rightarrow y_T$ of F , whose image by h is f such that: $l^{f'f} = l^{f'} \circ l^f$.

Consider the diagram

$$\begin{array}{ccc} F & & G \\ \downarrow f & & \downarrow g \\ E_1 & \xrightarrow{u} & E_2 \end{array}$$

where the functors f and g are fibred categories. We denote by $Hom_u(F, G)$ the subcategory of $Hom(F, G)$ whose objects are functors $v : F \rightarrow G$ which verify $gv = uf$. The maps of this category are morphisms $m : v \rightarrow v'$ such that gm is the identity morphism of the functor uf .

We denote by $Cart_u(F, G)$, the subcategory of $Hom_u(F, G)$ whose objects are cartesian functors: These are functors which transform cartesian maps to cartesian maps •

Let E be a category endowed with a topology J , and $F \rightarrow E$ a fibred category, for each object U of E , and each element R of $J(U)$, we consider the canonical functors $E_U \rightarrow E$, and $R \rightarrow E$. We can define the set of cartesian functors $Cart_{Id_E}(E_U, F)$ and $Cart_{Id_E}(R, F)$. There exists a canonical restriction functor $Cart_{Id_E}(E_U, F) \rightarrow Cart_{Id_E}(R, F)$.

Definition 2.0.6.

Let E be a category endowed with a topology, a **sheaf of categories** on E , is a fibred category $F \rightarrow E$, such that for each sieve R , the cartesian functor $Cart_{Id_E}(E_U, F) \rightarrow Cart_{Id_E}(R, F)$ defined at the paragraph above is an equivalence of categories •

Proposition-Definition 2.0.7.

Suppose that E is a topos whose topology is generated by a contractible covering family $(U_i \rightarrow U)_{i \in I}$, and $h : F \rightarrow E$ a fibred category in groupoids. For each map $f : U \rightarrow V$ of E , we consider the functor $r_{U,V}(f) : F_V \rightarrow F_U$ defined as follows: For each object y of F_V , $r_{U,V}(f)(y)$ is an object x of F_U such that there exists a cartesian map $n : x \rightarrow y$ such that $h(n) = f$. Consider the

maps $v_1 : U_1 \rightarrow U_2$, and $v_2 : U_2 \rightarrow U_3$ of E , the functors $r_{U_1, U_2}(v_1) \circ r_{U_2, U_3}(v_2)$ and $r_{U_1, U_3}(v_2 v_1)$ are isomorphic (see [11]). The functor $h : F \rightarrow E$ is a sheaf of categories if and only if the correspondence $U \rightarrow F_U = F(U)$ satisfies the following properties:

(i) *Gluing condition for arrows.*

Let U be an object of E , and x, y objects of $F(U)$. The functor from E_U , endowed with the restriction of the topology J , to the category of sets which associates to an object $f : V \rightarrow U$ the set $\text{Hom}_V(r_{V, U}(f)(x), r_{V, U}(f)(y))$ is a sheaf of sets.

(ii) *Gluing condition for objects.*

Consider a covering family $(U_i \rightarrow U)_{i \in I}$ of an object U of E , and for each U_i , an object x_i of $F(U_i)$. Let $t_{ij} : x_j^i \rightarrow x_i^j$, a map between the respective restrictions of x_j and x_i to $U_i \times_U U_j$ such that on $U_{i_1} \times_U U_{i_2} \times_U U_{i_3}$, the restrictions of the arrows $t_{i_1 i_3}$ and $t_{i_1 i_2} t_{i_2 i_3}$ are equal. There exists an object x of $F(U)$ whose restriction to $F(U_i)$ is x_i .

If moreover the following properties are verified:

(iii) There exists a covering family $(U_i \rightarrow U)_{i \in I}$ of E such that $F(U_i)$ is not empty,

(iv) For each pair of objects x , and y of $F(U_i)$, $\text{Hom}_{U_i}(x, y)$ is not empty (local connectivity),

(v) The elements of $\text{Hom}_{U_i}(x, y)$ are invertible. The fibered category is called a **gerbe**.

(vi) We say that the gerbe is **bounded** by the sheaf L_F defined on E , or that L_F is the **band** of the gerbe, if and only if there exists a sheaf of groups L_F defined on E such that for each object x of $F(U)$ we have an isomorphism:

$$L_F(U) \rightarrow \text{Hom}_U(x, x)$$

which commutes with restrictions, and with morphisms between objects •

2.1 Classifying cocycle and classification of gerbes.

In this paragraph, we recall the definition of the classifying cocycle of a gerbe defined on the topos E and bounded by the sheaf L .

Definitions 2.1.1.

- A gerbe $F \rightarrow E$ is trivial if it has a section. This means that F_E is not empty.
- Two gerbes $F \rightarrow E$, and $F' \rightarrow E$ whose band is L , are equivalent if and only if there exists an isomorphism between the underlying fibered categories which commutes with the action of L . We denote by $H^2(E, L)$ the set of equivalence classes of gerbes defined on E bounded by L •

Suppose that the topology of E is defined by the covering family $(U_i \rightarrow U)_{i \in I}$, the class of objects of F_{U_i} is not empty, and each objects x and y of F_{U_i} are isomorphic. Let $(x_i)_{i \in I}$ be a family of objects of F , such that x_i is an object of F_{U_i} . There exists a map $u_{ij} : x_j^i \rightarrow x_i^j$ between the respective restrictions of x_j and x_i to $F_{U_i \times_U U_j}$. We denote by $u_{i_1 i_2}^{i_3}$ the map between the respective restrictions of x_{i_2} and x_{i_1} to $F_{U_{i_1} \times_U U_{i_2} \times_U U_{i_3}}$.

Theorem [11] 2.1.2.

The family of maps $c_{i_1 i_2 i_3} = u_{i_3 i_1}^{i_2} u_{i_1 i_2}^{i_3} u_{i_2 i_3}^{i_1}$ is the classifying 2-Cech cocycle of the gerbe. If the band L is commutative, then the set of equivalence classes of gerbes over E whose band is L , is one to one with the Čech cohomology group $H^2(E, L)$.

3 Notations.

Let $U_{i_1}, \dots, U_{i_p}$ be objects of a topos E , and C a presheaf of categories defined on E . We will denote by $U_{i_1 \dots i_p}$ the fiber product of $U_{i_1}, \dots, U_{i_p}$ over the final object. If e_{i_1} is an object of $C(U_{i_1})$, $e_{i_1}^{i_2 \dots i_p}$ will be the restriction of e_{i_1} to $U_{i_1 \dots i_p}$. For a map $h : e \rightarrow e'$ between two objects of $C(U_{i_1 \dots i_p})$, we denote by $h^{i_{p+1} \dots i_n}$ the restriction of h to a morphism between $e^{i_{p+1} \dots i_n} \rightarrow e'^{i_{p+1} \dots i_n}$.

4 Gerbed tower.

The purpose of this part is to generalize the notion of gerbe to the notion of gerbed tower. This notion will allow us to define, and to represent geometrically higher non abelian cohomological classes. In the sequel we assume known the notion of 2-category or bicategory defined by Benabou [1]. Recall that a 2-category C is defined by a class of objects $Obj(C)$, and for each objects x and y of C , a category $Hom_C(x, y)$ called the category of morphisms. The objects of $Hom_C(x, y)$ are called 1-arrows, and the arrows are called 2-arrows. There exists a composition functor:

$$c(u_1, u_2, u_3) : Hom(u_2, u_3) \times Hom(u_1, u_2) \longrightarrow Hom(u_1, u_3)$$

For each quadruple (u_1, u_2, u_3, u_4) in C , there exists an isomorphism $c(u_1, u_2, u_3, u_4)$ between the functors

$$(Hom(u_3, u_4) \times Hom(u_2, u_3)) \times Hom(u_1, u_2) \longrightarrow Hom(u_1, u_4)$$

and

$$Hom(u_3, u_4) \times (Hom(u_2, u_3) \times Hom(u_1, u_2)) \longrightarrow Hom(u_1, u_4)$$

which satisfies more compatibility axioms which can be found in Benabou. We will suppose that $c(u_1, u_2, u_3, u_4)$ is the identity on objects. This implies that we can define the category C_1 whose objects are the objects of C , and such that for each pair of objects x, y of C_1 , $Hom_{C_1}(x, y)$ is the set of objects of $Hom_C(x, y)$. Let $h : F \rightarrow E$ be a gerbe. We can define the 2-category $C(E, F)$ whose objects are objects of F . Let x and y be a pair of objects of $C(E, F)$, an object of $Hom_{C(E, F)}(x, y)$ is an arrow between $h(x)$ and $h(y)$. A 2-arrow between the objects x and y is a cartesian map between x and y .

Definition 4.0.1.

A **bicategory C , endowed with a topology J** , is a bicategory whose objects are toposes, and for each pair of objects x and y of C . The set of 2-arrows between x and y is contained in the space of continuous maps between x and y •

Definition 4.0.2.

A **n -gerbed tower** is defined by:

1. A family $F_n, F_{n-1}, \dots, F_2, F_1$ of 2-categories respectively endowed with topologies $J_n, \dots, J_1$, and a family of 2-functors $p_l : F_l \rightarrow F_{l-1}$, $l \in \{2, \dots, n\}$ which satisfy the following conditions:

2. F_1 is a gerbe $p_1 : F \rightarrow E$, since we can assume that E is a 2-category such that for each objects U and V of E , the set of arrows between a pair of elements f and f' of $Hom_E(U, V)$ is a singleton, we will often consider the sequence of 2-categories $F_n \rightarrow \dots F_1 \rightarrow E$. We suppose that the 2-arrows of F_p , $p \in \{1, \dots, n\}$ are invertible.

3. Let U be an object of F_p , and $l \geq p$ a pair of integers inferior to n . We denote by F_{lpU} , the 2-category whose class of objects is contained in the class of objects of F_l , and such that V is an object of F_{lpU} if and only if $p_{p+1} \dots p_l(V) = U$. The category of morphisms $Hom_{F_{lpU}}(X, Y)$ between a pair of objects X and Y is the subcategory of $Hom_{F_l}(X, Y)$ such that the projections of 1-arrows of $Hom_{F_{lpU}}(X, Y)$ by $p_{p+1} \dots p_l$ is Id_U of U , and the projections of 2-arrows of $Hom_{F_{lpU}}(X, Y)$ by the same functor is the identity of Id_U . We denote F_{l0U} by F_{lU} . We suppose that for each arrow $f : U \rightarrow V$ of E , there exists a restriction functor $r_{U,V}^l(f) : F_{lV} \rightarrow F_{lU}$ such that for every map $g : V \rightarrow V'$, $r_{U,V}^l(f) \circ r_{V,V'}^l(g) = r_{U,V'}^l(gf)$.

4. There exists a family of sheaves $L_1, \dots, L_n$ defined on E . The sheaf L_{l+1} induces a sheaf L_{l+1U_l} on the object U_l of F_l (recall that U_l is a topos) defined by its global sections $L_{l+1U_l} = L_{l+1}(p_1 \dots p_l(U_l))$. For each object U_{l-1} of F_{l-1} , we suppose that the fiber of $F_{ll-1U_{l-1}}$ is a gerbe defined on the topos U_{l-1} bounded by $L_{lU_{l-1}}$.

5. Let U_{1l} and U_{2l} be a pair of objects of F_l , and $u_l^2 : h_l^1 \rightarrow h_l^2$ a 2-arrow, between U_{1l} and U_{2l} , that is an arrow of the category $Hom_{F_l}(U_{1l}, U_{2l})$ between the objects h_l^1 and h_l^2 . Recall that u_l^2 is a continuous functor between the topoi U_{1l} and U_{2l} . We suppose that for every object U'_{2l} of $F_{l+1lU_{2l}}$ there exists an object U'_{1l} of $F_{l+1lU_{1l}}$, and a 2-arrow $u_l^{2*} : U'_{1l} \rightarrow U'_{2l}$ such that the following diagram is commutative:

$$\begin{array}{ccc} U'_{1l} & \xrightarrow{u_l^{2*}} & U'_{2l} \\ \downarrow p_l & & \downarrow p_l \\ U_{1l} & \xrightarrow{u_l^2} & U_{2l} \end{array}$$

This implies that for every 2-arrow $v_l^2 : h_l^2 \rightarrow h_l^3$ between the pair of objects U_{2l} and U_{3l} , for every object U'_{3l} of $F_{l+1lU_{3l}}$, and every 2-arrow v_l^{2*} of F_{l+1} defined by the diagram above, there exists an automorphism over the identity $c(u_l^2, v_l^2)$ of U'_{3l} such that

$$v_l^{2*} \circ u_l^{2*} = c(u_l^2, v_l^2)(v_l^2 \circ u_l^2)^*,$$

since we have supposed that the 2-arrows are invertible morphisms of topoi $\bullet$

Definition 4.0.3.

An ∞ -gerbed tower, is a sequence of functors between 2-categories $\dots F_n \rightarrow F_{n-1} \dots \rightarrow F_1 \rightarrow E$ such that for each integer n , $F_n \rightarrow F_{n-1} \dots \rightarrow F_1 \rightarrow E$ is a gerbed tower $\bullet$

Definitions 4.0.4.

- A morphism F between the gerbed towers $f = F_n \rightarrow \dots F_1 \rightarrow E$ and $f' = F'_n \rightarrow \dots \rightarrow F'_1 \rightarrow E$, is defined by a family of 2-functors $f_l : F_l \rightarrow F'_l$ such that for each l , the following diagram is commutative:

$$\begin{array}{ccc} F_l & \xrightarrow{f_l} & F'_l \\ \downarrow p_l & & \downarrow p'_l \\ F_{l-1} & \xrightarrow{f_{l-1}} & F'_{l-1} \end{array}$$

and for each object U_l of F_l , the induced morphism: $F_{l+1}U_l \rightarrow F'_{l+1}f_l(U_l)$ is a morphism of gerbes.

- The morphism defined by the family of 2-functors $(f_n, \dots, f_1)$ is an isomorphism, if and only if there exists a morphism between the gerbed tower $F'_n \rightarrow \dots \rightarrow F'_1 \rightarrow E$ and $F_n \rightarrow \dots \rightarrow F_1 \rightarrow E$ defined by the family of 2-functors $f'_n, \dots, f'_1$ such that for each l , $f'_l \circ f_l = Id_{F_l}$, and $f_l \circ f'_l = Id_{F'_l}$.

We say that the gerbed towers f' and f are weakly equivalent if and only if $f'_l \circ f_l$ is isomorphic to Id_{F_l} , and $f_l \circ f'_l$ is isomorphic to $Id_{F'_l}$. We denote by $H^{n+1}(E, L_n)$ the set of weakly equivalence classes of gerbed towers bounded by $(L_1, \dots, L_n)$. Here L_n is a fixed sheaf defined on E •

4.1 Non commutative cohomology of groups.

Let H be a group, V a vector space, $Gl(V)$ the group of linear automorphisms of V , and $\rho : H \rightarrow Gl(V)$ a representation. To define the cohomology groups $H^n(H, V, \rho)$ of the representation ρ , one can consider EH the 1-Eilenberg-MacLane space defined by H , the representation ρ defines on EH a flat V -bundle whose holonomy is ρ . The cohomology groups $H^n(H, V, \rho)$, are the n -cohomology groups, of the sheaf of locally constant sections of this flat bundle. This motivates the following definition:

Definition 4.1.1.

Consider the groups H and G , $Aut(G)$ the group of automorphisms of G , and $\rho : H \rightarrow Aut(G)$ a representation. The representation ρ defines on EH a flat G -bundle p_G . We denote by L_{p_G} the sheaf of locally constant sections of this bundle. We define $H^{n+1}(H, G, \rho)$ to be set of weakly equivalence classes of gerbed towers of rank n $F_n \rightarrow \dots \rightarrow F_1 \rightarrow EH$ bounded by a sequence $(L_1, \dots, L_{n-1}, L_{p_G})$ •

4.2 The classifying cocycle of a gerbed tower.

Let $f = F_n \rightarrow F_{n-1} \rightarrow \dots \rightarrow F_1 \rightarrow E$ be a gerbed tower. We will define in this part the classifying cocycle of f . We suppose that the sheaves $L_1, \dots, L_n$ are commutative, and there exists commutative sheaves $L'_1, \dots, L'_n$ defined on E such that for every objects U_l of F_l , U_{l+1} of $F_{l+1}U_l$, and h_l an object of $Hom_{F_{l+1}}(U_{l+1}, U_{l+1})$, $Aut(F_{l+1}h_{l+1}) = L'_{l+1}(p_1 \dots p_l(U_l))$ where $Aut(F_{l+1}h_{l+1})$ is the group of automorphisms of the object h_{l+1} whose image by p_{l+1} are elements of $L_l(p_1 \dots p_l(U_l))$.

Suppose that the topology of E is defined by the covering family $(U_i \rightarrow U)_{i \in I}$ such that for each i , F_{1U_i} is not empty, and its objects are isomorphic. Let u_i be an object of F_{1U_i} , and $v_{i_1 i_2}$ an arrow between $u_{i_2}^{i_1}$ and $u_{i_1}^{i_2}$. The family of arrows $c_{i_1 i_2 i_3} = v_{i_3 i_1}^{i_2} v_{i_1 i_2}^{i_3} v_{i_2 i_3}^{i_1}$ is the classifying cocycle of the gerbe $F_1 \rightarrow E$.

If we identify $F_1 \rightarrow E$ with a 2-category, then $c_{i_1 i_2 i_3}$ is a 2-arrow. Let $u_{i_1 i_2 i_3}$ be an object of the fiber $F_{21}u_{i_3}^{i_1 i_2}$. The property 5 of the definition of gerbed towers implies the existence of a 2-arrow $c_{i_1 i_2 i_3}^*$ of $Hom_{F_2}(u_{i_1 i_2 i_3}, u_{i_1 i_2 i_3})$ over $c_{i_1 i_2 i_3}$.

we can define the automorphism $c_{i_1 i_2 i_3 i_4}$ of the object $u_{i_1 i_2 i_3}^{i_4}$ by:

$$c_{i_1 i_2 i_3 i_4} = c_{i_2 i_3 i_4}^{*i_1} - c_{i_1 i_3 i_4}^{*i_2} + c_{i_1 i_2 i_4}^{*i_3} - c_{i_1 i_2 i_3}^{*i_4}$$

each member of the right part of the previous equality can be supposed to be a morphism of the same object of $\text{Hom}_{F_2}(u_{i_1 i_2 i_3}, u_{i_1 i_2 i_3})$. The property 5 of the definition of gerbed towers implies that $c_{i_1 i_2 i_3 i_4}$ is an element of $L_2(U_{i_1} \times_E U_{i_2} \times_E U_{i_3} \times_E U_{i_4})$.

Proposition 4.2.1.

The family $c_{i_1 i_2 i_3 i_4}$ that we have just defined is a 2-Cech cocycle.

Proof.

The Cech boundary of $c_{i_1 i_2 i_3 i_4}$ is:

$$\begin{aligned} \partial(c_{i_1 \dots i_4}) &= \sum_{p=1}^{p=5} (-1)^j c_{i_1 \dots \hat{i}_p \dots i_5} \\ &= \sum_{p=1}^{p=5} (-1)^p \left(\sum_{c=1}^{c=p-1} (-1)^c c_{i_1 \dots \hat{i}_c \dots \hat{i}_p \dots}^* + \sum_{c=p+1}^{c=5} (-1)^{c+1} c_{i_1 \dots \hat{i}_p \dots \hat{i}_c \dots i_5}^* \right) = 0. \end{aligned}$$

The last sum is zero because the sheaf L'_1 is commutative •

Suppose defined the classifying cocycles $c_{i_1 i_2 i_3}, \dots, c_{i_1 \dots i_{l+2}}$, $l \geq 2$ of the gerbed tower $F_l \rightarrow \dots \rightarrow F_1 \rightarrow F$. The arrow $c_{i_1 \dots i_{l+2}}$ is a 2-arrow of $u_{i_1 \dots i_{l+1}}^{i_{l+2}}$. Let $u_{i_1 \dots i_{l+2}}$ be an object of $F_{l+1} u_{i_1 \dots i_{l+1}}^{i_{l+2}}$. The property 5 implies the existence of an automorphism $c_{i_1 \dots i_{l+2}}^*$ of a 1-arrow of the object $u_{i_1 \dots i_{l+2}}$ over $c_{i_1 \dots i_{l+2}}$. We can define:

$$c_{i_1 \dots i_{l+3}} = \sum_{p=1}^{l+3} (-1)^p c_{i_1 \dots \hat{i}_p \dots i_{l+3}}^*$$

We can apply the property 5 to identify $c_{i_1 \dots i_{l+3}}$ to a 2-arrow of $u_{i_1 \dots i_{l+2}}^{i_{l+3}}$.

Proposition 4.2.2.

The family of arrows $c_{i_1 \dots i_{l+3}}$ that we have just defined is a $l+1$ -Cech cocycle.

Proof.

The Cech boundary of $c_{i_1 \dots i_{l+3}}$ is:

$$\begin{aligned} \partial(c_{i_1 \dots i_{l+3}}) &= \sum_{p=1}^{p=l+4} (-1)^p c_{i_1 \dots \hat{i}_p \dots i_{l+4}} \\ &= \sum_{p=1}^{p=l+4} (-1)^p \left(\sum_{c=1}^{c=p-1} (-1)^c c_{i_1 \dots \hat{i}_c \dots \hat{i}_p \dots}^* + \sum_{c=p+1}^{c=l+4} (-1)^{c+1} c_{i_1 \dots \hat{i}_p \dots \hat{i}_c \dots i_{l+4}}^* \right) = 0. \end{aligned}$$

The last sum is zero because the sheaf L'_{l+1} is commutative •

Proposition 4.2.3.

The cohomology class $[c_{l+2}]$, is the image of $[c_{l+1}]$ by the connecting morphism $H^{l+1}(E, L_l) \rightarrow H^{l+2}(E, L_{l+1})$ of the exact sequence $0 \rightarrow L_{l+1} \rightarrow L'_{l+1} \rightarrow L_l \rightarrow 0$. In particular this shows that the cohomology classes of the cocycles c_l , $2 \leq l \leq n+1$ attached to the gerbed tower $F_n \rightarrow F_{n-1} \dots \rightarrow F_1 \rightarrow E$ are independent of the choices made to construct them.

Proof.

To construct the cocycle c_{l+2} we pick elements $c_{i_1 \dots i_{l+2}}$ which represents 2-arrows of an object $u_{i_1 \dots i_{l+1}}^{i_{l+2}}$ that we lift to 2-arrows $c_{i_1 \dots i_{l+2}}^*$ of the object $u_{i_1 \dots i_{l+2}}$ of $F_{l-1} u_{i_1 \dots i_{l+1}}^{i_{l+2}}$. The representant $c_{i_1 \dots i_{l+3}}$ of $[c_{l+2}]$ are the Cech boundary of the family $c_{i_1 \dots i_{l+2}}^*$ acting on $u_{i_1 \dots i_{l+2}}$. This is by definition the construction of the connecting morphism $H^{l+1}(E, L_l) \rightarrow H^{l+2}(E, L_{l+1})$ of the exact sequence $0 \rightarrow L_{l+1} \rightarrow L'_{l+1} \rightarrow L_l \rightarrow 0$ •

Definition 4.2.4.

A gerbed tower $f_n = F_n \rightarrow F_{n-1} \dots F_1 \rightarrow E$ is trivial if and only if there exists a gerbed tower $f_{n-1} = F'_{n-1} \rightarrow F'_{n-2} \dots F'_1 \rightarrow E$ and an element $p \in \{1, \dots, n-2\}$ such that F'_l is F_l if $l \leq p$, F'_p is a sub 2-category of F_{p+1} , for each 2-arrow h of F_{p-1} , the image of the arrow h^* of F'_p by p_{p+1} is an arrow u^* , of F_p , defined by the axiom 5, defined by a 2-arrow of u of F_p . For $l > p$, the 2-category F'_l is a subcategory of F_{l+1} . The gerbed tower f_{n-1} is called a trivialization of f_n •

Proposition 4.2.5.

The class $[c_{n+1}]$ of a trivial gerbed tower $F_n \rightarrow \dots \rightarrow F_1 \rightarrow E$ is zero.

Proof.

Let $f_n = F_n \rightarrow F_{n-1} \rightarrow \dots F_1 \rightarrow E$ be a trivial gerbed tower, and $F'_{n-1} \rightarrow F'_{n-2} \dots \rightarrow F'_1 \rightarrow E$ a trivialization of f_n . Suppose that the integer p of the definition above is $n-2$, this means that if $l \leq n-2$, and F'_l is F_l . We denote by $(L_1, \dots, L_n)$ the band of the gerbed tower $F_n \rightarrow \dots F_1 \rightarrow E$, and by L''_n the sheaf such that the group $\text{Aut}(F'_{n-1} h_{n-1})$ of automorphisms of a 2-arrow h_{n-1} of the object U_{n-1} of F'_{n-1} which project by $p'_1 \dots p'_{n-1}$ to elements of $L_{n-2}(p'_1 \dots p'_{n-1}(U_{n-1}))$ is $L''_n(p_1 \dots p'_{n-1}(U_{n-1}))$. We have the commutative:

$$\begin{array}{ccccccc} 0 & \longrightarrow & L_n & \longrightarrow & L'_n & \longrightarrow & L_{n-1} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & L_n & \longrightarrow & L''_n & \longrightarrow & L_{n-2} \longrightarrow 0 \end{array}$$

The map of $L'_n \rightarrow L''_n$ is defined by the restriction of the morphisms u^* , where $u \in L_{n-1}$, and the map $L_{n-1} \rightarrow L_{n-2}$ is zero. This exact sequence gives rise to the commutative diagram:

$$\begin{array}{ccc} H^n(E, L_{n-1}) & \longrightarrow & H^n(E, L_{n-2}) \\ \downarrow & & \downarrow \\ H^{n+1}(E, L_n) & \longrightarrow & H^{n+1}(E, L_n) \end{array}$$

Since the map $H^n(E, L_{n-1}) \rightarrow H^n(E, L_{n-2})$ is zero, and the map $H^{n+1}(E, L_n) \rightarrow H^{n+1}(E, L_n)$ is the identity, we deduce that the class $[c_{n+1}]$ of the classifying cocycle of the gerbed tower $F_n \rightarrow F_{n-1} \dots \rightarrow F_1 \rightarrow E$ is zero.

If p is not $n-1$, the last argument show that $[c_{p+1}]$ is zero. This implies that $[c_l] = 0$ for $l \geq p+1$ •

Theorem 4.2.6.

Let $F_n \rightarrow F_{n-1} \dots F_1 \rightarrow E$ be a gerbed tower bounded by the family of sheaves $(L_1, \dots, L_n)$ whose classifying cocycles are $(c_2, \dots, c_{n+1})$. Consider an exact sequence of sheaves $0 \rightarrow L_{n+1} \rightarrow L'_{n+1} \rightarrow L_n \rightarrow 0$. Then there exists a gerbed tower $F_{n+1} \rightarrow F_n \dots \rightarrow F_1 \rightarrow E$ whose classifying cocycles are $c_2, \dots, c_{n+1}, c_{n+2}$, where c_{n+2} is a $n+1$ -cocycle whose cohomology class is the image of the class of c_{n+1} by the connecting map $H^{n+1}(E, L_n) \rightarrow H^{n+2}(E, L_{n+1})$ defined by the previous exact sequence.

Proof.

Let U_n be an object of F_n , and U_{n-1} its image by the projection map $p_n : F_n \rightarrow F_{n-1}$. The topos U_n is a L_n -torsor defined over an object of the topos U_{n-1} since $F_{nn-1}U_{n-1}$ is a L_n -gerbe defined on U_{n-1} . An object V_n of $F_{nn-1}U_{n-1}$ is a L_n -torsor defined over an object U'_{n-1} of the topos U_{n-1} . This torsor is defined by a trivialization $(V_i)_{i \in I}$, and coordinate changes $u_{ij} : V_i \times_{U_{n-1}} V_j \rightarrow L_n$. This coordinate changes define a principal L_n -torsor over U'_{n-1} by gluing $V_i \times L_n$. Without restricting the generality, we can suppose that the objects of $F_{nn-1}U_{n-1}$ are L_n -principal torsors. We define the fiber $F_{n+1}U_n$ to be the gerbe bounded by L_{n+1} defined on U_n which represents the obstruction to lifts the L_n -torsor $p_{U_n} : U_n \rightarrow U'_{n-1}$ to a L'_{n+1} -torsor whose quotient by L_{n+1} is the previous L_n -torsor $p_{U_n} : U_n \rightarrow U'_{n-1}$.

The objects of the category of morphisms $Hom_{F_{n+1}}(U_{n+1}, U'_{n+1})$ between the objects U_{n+1} of $F_{n+1}U_n$ and U'_{n+1} of $F_{n+1}U'_n$ are the 2-arrows between U_n and U'_n of F_n , the 2-arrows are the morphisms of torsors u^2_{n+1} such that there exists a 2-arrow $u^1_n : U_n \rightarrow U'_n$ such that the following diagram is commutative:

$$\begin{array}{ccc} U_{n+1} & \xrightarrow{u^2_n} & U'_{n+1} \\ \downarrow p_{n+1} & & \downarrow p_{n+1} \\ U_n & \xrightarrow{u^1_n} & U'_n \end{array}$$

We show now that $F_{n+1} \rightarrow F_n \dots \rightarrow F_1 \rightarrow E$ is a gerbed tower:

Let U_n be an object of F_n , and U the object $p_1 \dots p_n(U_n)$ of E . For each map $f : V \rightarrow U$, we can define the restriction $r^n_{V,U}(f) : F_{nU} \rightarrow F_{nV}$. The restriction $r^{n+1}_{V,U}$ is defined on $F_{n+1}U_n$ by the pull-back of U_{n+1} by the arrow $r^n_{V,U}(f)$.

The definition of F_{n+1} implies that for every object U_n of F_n , the gerbe $F_{n+1}U_n$ is bounded by L_{n+1} .

For every 2-arrow $u^2_n : h^1_n \rightarrow h^2_n$, (recall that u^2_n is a morphism between the topoi U_{1n} and U_{2n}) of F_n , the functor u^{2*}_n is defined as follows: Without restricting the generality, we can suppose that U_{1n} is the trivial torsor to $V_1 \times L_n$ and U_{2n} the trivial torsor $V_2 \times L_n$; u^2_n is then a morphism of L_n -torsors, u^{2*}_n is a morphism such that the following diagram is commutative:

$$\begin{array}{ccc} V_1 \times L'_{n+1} & \xrightarrow{u^{2*}_n} & V_2 \times L'_{n+1} \\ \downarrow & & \downarrow \\ V_1 \times L_n & \xrightarrow{u^2_n} & V_2 \times L_n \end{array}$$

This shows that $F_{n+1} \rightarrow \dots F_1 \rightarrow E$ is a gerbed tower.

The classifying cocycle of this gerbed tower is constructed by considering the automorphism $c_{i_1 \dots i_{n+2}}$ of the object $u_{i_1 \dots i_{n+2}}$ of F_{n+1} , that we suppose to be isomorphic to a trivial torsor $V_{i_1 \dots i_{n+2}} \times L_n$, the morphism $c_{i_1 \dots i_{n+2}}$ can be lifted to an element $c_{i_1 \dots i_{n+2}}^*$ of $V_{i_1 \dots i_{n+2}} \times L'_{n+1}$. The Čech boundary of $c_{i_1 \dots i_{n+2}}^*$ is the classifying cocycle of the gerbed tower. The cohomology class of this cocycle is the image of the cohomology class of c_{n+1} by the connecting morphism $H^{n+1}(E, L_n) \rightarrow H^{n+2}(E, L_{n+1})$ •

5 Spectral sequences and gerbed towers.

The goal of this part is to apply spectral sequences to study commutative gerbed towers.

Let $E(L_1, \dots, L_n, \dots)$ be an ∞ -gerbed tower, where $(L_n)_{n \in \mathbb{N}}$ is a family of commutative sheaves defined on E . We suppose that the topology of E is defined by the covering family $(X_i \rightarrow X)_{i \in I}$. We define $L = \bigoplus_{i \geq 1} L_i$, and denote by $(C^*(X, L), d)$ the complex of Čech L -chains defined on E . We can endow this chain complex with the following filtration:

$$V_p = C(X, \bigoplus_{i \geq p} L_i),$$

and with the graduation

$$V^p = C^p(X, L)$$

We will calculate the terms associated to the spectral sequence associated to this graduation.

Denote by $Z_r^p = \{x \in V_p : d(x) \in V_{p+r}\}$, $B_r^p = d(V_{p-r}) \cap V_p$, and $E_r^p = \frac{Z_r^p}{Z_{r-1}^{p+1} + B_{r-1}^p}$.

We suppose in the sequel that $r \geq 1$.

Determination of Z_r^p .

Let x be an element of V_p , $d(x)$ is an element of Z_r^p if and only if $d(x)$ is an element of V_{p+r} . We can write $x = x_{i_p} + \dots + x_{i_n}$, where x_{i_l} is the homogeneous component of x which takes value in L_l , $d(x_l)$ is an element of V_{p+r} if and only if $d(x_l) = 0$ if $l \leq p+r$. We deduce that x is an element of Z_r^p if and only if its components x_{i_j} , such that $j < p+r$ are cocycles.

Determination of B_r^p .

Let x be an element of V_{p-r} , and x_l its component which takes values in L_l . The image by d of x_l is an element of B_r^p if and only if $d(x_l)$ is an element of V_p . This equivalent to saying that $d(x_l)$ is zero, or $l \geq p$. This implies that $B_r^p = d(V_p)$.

Determination of E_r^p .

We have $Z_r^p = Z_{r-1}^{p+1} \oplus Z(V_p) \cap C(X, L_p)$. We deduce that $E_r^p = H(X, L_p)$.

Now we set $Z_r^{pq} = Z_r^p \cap V^{p+q}$, $B_r^{pq} = B_r^p \cap V^{p+q}$ and $E_r^{pq} = \frac{Z_r^{pq}}{B_{r-1}^{pq} + Z_{r-1}^{p+1, q-1}}$.

Determination of Z_r^{pq} .

Let x be an element of Z_r^p , one of its homogeneous components x_l which takes values in L_l , is an element of Z_r^{pq} , if x_l is a V_p $p+q$ -chain, and $d(x_l)$ is an element of V_{p+r} . We have seen that $d(x_l)$ is an element of Z_r^p if and only if x_l is a cocycle or $r \geq p+r$. We deduce from this fact that $Z_r^{pq} = C^{p+q}(X, V_{p+r}) \oplus Z^{p+q}(X, L_1 \oplus \dots \oplus L_{p+r-1})$.

Determination of B_r^{pq} .

Let x be an element of B_r^{pq} , one of its homogeneous components, x_l which takes value in L_l is an element of B_r^{pq} if and only if it is a $p+q$ -chain, and there exists an element y in V_{p-r} such that $d(y) = x_l$. We deduce that $B_r^{pq} = d(C^{p+q-1}(X, V_p))$.

Determination of E_r^{pq} .

The vector space Z_r^{pq} is the summand of $Z_{r-1}^{p+1,q-1}$ and $Z^{p+q}(X, L_p)$. We deduce that $E_r^{pq} = H^{p+q}(X, L_p)$.

Now, we will denote by Z_∞^p , the set of cocycles contained in V_p , by B_∞^p the set of boundaries contained in V_p , and by $E_\infty^p = \frac{Z_\infty^p}{Z_\infty^{p+1} + B_\infty^p}$. We remark that $E_\infty^p = H(X, L_p)$.

The following proposition can be deduced from [19] p. 84 Theorem 4.6.1.

Proposition 4.0.1.

Suppose that there exists an integer $n \geq r$ such that $H^{p+q}(X, L_p) = E_r^{pq} = 0$ for $p \neq 0, n$ and an integer s such that $L_n = 0$ if $n > s$, then we have the following exact sequence

$$\dots \rightarrow H^i(X, L_n) \rightarrow H^i(X, L) \rightarrow H^i(X, L_1) \rightarrow H^{i+1}(X, L_n) \rightarrow H^{i+1}(X, L) \rightarrow \dots$$

6 Application of gerbed towers to affine manifolds.

An affine manifold (N, ∇_N) , is a differentiable manifold N , endowed with a connection ∇_N whose curvature and torsion forms vanish identically. The connection ∇_N defines on N an atlas whose coordinate changes are affine transformations. Auslander has conjectured that the fundamental group of a compact and complete affine manifold is polycyclic. Let (N, ∇_N) and $(N', \nabla_{N'})$ be two affine manifolds of respective dimension n and n' whose affine structures are defined by the respective atlases (U_i, u_i) , and (U'_j, u'_j) . An affine map $f : (N, \nabla_N) \rightarrow (N', \nabla_{N'})$ is a differentiable map $f : N \rightarrow N'$ such that $u'_i \circ f|_{U_i} \circ u_i^{-1}$ is a restriction of an affine map from $\mathbb{R}^n$ to $\mathbb{R}^{n'}$. Suppose that (N, ∇_N) and $(N', \nabla_{N'})$ are complete and compact. It is shown in Tsemo [26] that in this case, there exists a compact and complete affine manifold (N_1, ∇_{N_1}) of dimension n' , and an affine submersion $f_1 : (N, \nabla_N) \rightarrow (N_1, \nabla_{N_1})$. Ehresman has shown that submersions between compact manifolds are locally trivial differentiable fibrations. The typical fiber F of the fibration $N \rightarrow N_1$ inherits from N complete affine structures. The homotopy exact sequence of this fibration gives rise to the sequence:

$$1 \longrightarrow \pi_1(F) \rightarrow \pi_1(N) \rightarrow \pi_1(N_1) \rightarrow 1$$

If the fundamental groups of $\pi_1(F)$ and $\pi_1(N_1)$ are polycyclic, then $\pi_1(N)$ is also polycyclic. This has motivated the following conjecture:

Conjecture 5.0.1.

Let (N, ∇_N) be a n -dimensional compact and complete affine manifold, then there exists a complete affine structure $(N', \nabla_{N'})$ defined on a finite galois cover N' of N , and a non trivial affine map $f : (N', \nabla_{N'}) \rightarrow (N_1, \nabla_{N_1})$. Non trivial means that the dimension of the fibers of f are different from zero, and n .

This conjecture implies the Auslander conjecture, and leads to the problem of classifying sequences of affine submersions $(N_n, \nabla_{N_n}) \rightarrow \dots \rightarrow (N_1, \nabla_{N_1})$. This if the last conjecture is true, will allow to know the topology of all the compact and complete affine manifolds. The theory of gerbed towers has been first constructed to study this classification problem.

Definition 5.0.2.

An affinely locally trivial affine fibration, whose typical fiber is the affine manifold (F, ∇_F) , is an affine map $f : (N_1, \nabla_{N_1}) \rightarrow (N, \nabla_N)$ which is the total space of a bundle whose fibers inherit from (N_1, ∇_{N_1}) , affine structures whose holonomies is the holonomy of the affine structure (F, ∇_F)

•

We will restrict to the study of sequences $(N_n, \nabla_{N_n}) \rightarrow \dots (N_1, \nabla_{N_1}) \rightarrow (N, \nabla_N)$ where each map $f_p : (N_p, \nabla_{N_p}) \rightarrow (N_{p-1}, \nabla_{N_{p-1}})$ is an affinely locally trivial affine fibration.

Let $f : (N_1, \nabla_{N_1}) \rightarrow (N, \nabla_N)$ be an affinely locally trivial affine fibration whose typical fiber is the affine manifold (F, ∇_F) . We suppose that the affine structure of (N_1, ∇_{N_1}) is complete. This implies that the affine structure of (N, ∇_N) is complete see Tsemo [27]. We can identify $\pi_1(N_1)$ with its image by the holonomy morphism of (N_1, ∇_{N_1}) . Suppose that the dimension of N_1 and N are respectively n_1 and n . Let $h : \pi_1(N_1) \rightarrow \text{Aff}(\mathbb{R}^{n_1})$ be the holonomy representation of (N_1, ∇_{N_1}) . We can write see Tsemo [27] $\mathbb{R}^{n_1} = \mathbb{R}^n \times \mathbb{R}^p$, and for each element γ of $\pi_1(N_1)$, $\gamma(x, y) = (L_{1\gamma}(x) + l_{1\gamma}, L_{2\gamma}(y) + L_{3\gamma}(x) + l_{2\gamma})$. Where $L_{1\gamma}$ and $L_{2\gamma}$ are respective automorphisms of $\mathbb{R}^n$ and $\mathbb{R}^p$, $L_3 : \mathbb{R}^n \rightarrow \mathbb{R}^p$ is a linear map, and $l_{1\gamma}$ and $l_{2\gamma}$ are respective elements of $\mathbb{R}^n$ and $\mathbb{R}^p$. An element γ of $\pi_1(F)$ is an element γ of $\pi_1(N_1)$ such that $L_{1\gamma}$ is the identity of $\mathbb{R}^n$, and $l_{1\gamma}$ is zero, and $L_{3\gamma}$ is zero. We can identify $\pi_1(N)$ to the set of affine transformations $(L_{1\gamma}, l_{1\gamma})$.

Let T_F be the translation group of (F, ∇_F) , that is the group of affine automorphisms of (F, ∇_F) whose elements lift to translations of $\mathbb{R}^p$. Since the group $\pi_1(F)$ is a normal subgroup of $\pi_1(N_1)$, the holonomy of (N_1, ∇_{N_1}) induces a representation $\pi_1(N) \rightarrow \text{Aff}(F, \nabla_F)/T_F$, which defines a flat $\text{Aff}(F, \nabla_F)/T_F$ -bundle p_F over (N, ∇_N) . The composition of the holonomy of (N_1, ∇_{N_1}) , and the conjugation of $\text{Aff}(F, \nabla_F)$ defined a flat bundle T_F -bundle p'_F over (N, ∇_N) see Tsemo [27]. An isomorphism $h : e \rightarrow e'$ between a pair of locally trivial (F, ∇_F) -affine bundles e and e' defined over (N, ∇_N) , is an affine map $h : e \rightarrow e'$ which is an isomorphism of bundles which gives rise to the identity of p_F .

Given affine manifolds (F, ∇_F) and (N, ∇_N) and a flat $\text{Aff}(F, \nabla_F)/T_F$ -bundle p_F , we can define the first extension problem as follow: study the existence and classify affinely locally trivial affine fibrations $f : (N_1, \nabla_{N_1}) \rightarrow (N, \nabla_N)$ which give rise to the flat bundle p_F .

Proposition 5.0.3.

Let (F, ∇_F) and (N, ∇_N) be compact and complete affine manifolds, and p_F a flat $\text{Aff}(F, \nabla_F)/T_F$ -bundle defined on N , The bundle p_F induces a flat T_F -bundle p'_F . For each open subset U of N , define the category $C_F(U)$ to be the category whose objects are affinely locally trivial (F, ∇_F) -affine bundles which induce the restriction of p_F to U . A map $h : e_U \rightarrow e'_U$ between two objects of $C_F(U)$ is an isomorphism of affine bundles which gives rise to the identity of the restriction of p_F to U . The correspondence defined on the category of open subsets of N , by $U \rightarrow C_F(U)$, is a gerbe whose classifying cocycle represents the obstruction to the existence of an affinely locally trivial affine bundle, which gives rise to p_F . The gerbe $C_F(U)$ is a gerbe bounded by the sheaf of affine sections of p'_F , that we denote L_F .

Proof.

Gluing property for objects.

Let $(U_i)_{i \in I}$ be an open covering of an open subset U of N , e_i an object of $C_F(U_i)$, and $u_{ij} : e_j^i \rightarrow e_i^j$ an isomorphism such that $u_{i_1 i_2}^{i_3} u_{i_2 i_3}^{i_1} = u_{i_1 i_3}^{i_2}$. The definition of a bundle implies

the existence of a bundle e defined over U whose restriction to U_i is e_i . Since the coordinate changes u_{ij} are affine isomorphisms between affinely locally trivial (F, ∇_F) -affine bundles, this implies that e is an affinely locally trivial (F, ∇_F) -affine bundle.

Gluing conditions for arrows.

Let e and e' be two objects of $C_F(U)$, the correspondence defined on the open subsets of U by $V \rightarrow \text{Hom}(e|_V, e'|_V)$, where $e|_V$ and $e'|_V$ are the respective restrictions of e and e' to V is a sheaf of sets, since it is a sheaf of morphisms between two bundles.

This shows that C_F is a sheaf of categories. It remains to show that C is a gerbe.

Let $(U_i)_{i \in I}$ be an open covering of N by contractible open subsets which are domain of affine charts. Then $U_i \times (F, \nabla_F)$ is an object of $C_F(U_i)$.

Let U be an open subset of N , and e and e' a pair of objects of $C_F(U)$. The respective restrictions $e|_{U_i \cap U}$, and $e'|_{U_i \cap U}$ of e and e' to $U_i \cap U$ are isomorphic to $U_i \cap U \times (F, \nabla_F)$.

An isomorphism h , of an object e of $C_F(U)$, is an isomorphism of affine bundle which gives rise to the identity of the restriction of p_F to U . The restriction of h to $e|_{U \cap U_i}$ is an isomorphism h_i of the trivial bundle $U \cap U_i \times (F, \nabla_F)$. The fact that h gives rise to the identity of p_F , is equivalent to the fact that its restriction to a fiber yields to the identity of $\text{Aff}(F, \nabla_F)/T_F$. This implies that h_i is a T_F valued affine map, and h is a section of p'_F •

Proposition 5.0.4.

Suppose that the gerbe C_F is trivial, then the objects of $C_F(N)$ are diffeomorphic manifolds.

Proof.

Suppose that the gerbe C_F is trivial, then the holonomy of a global object (N_1, ∇_{N_1}) is defined by a representation $h_\gamma(x, y) = (L_{1\gamma}(x) + l_{1\gamma}, L_{2\gamma}(y) + l_{2\gamma})$ which defines an (F, ∇_F) -bundle. The objects of $C_F(N)$ are classified by $H^1(N, p'_F)$, the 1-cohomology group of the sheaf of affine sections of p'_F . An element of $H^1(N, p'_F)$ is defined by an affine $C_3 : \mathbb{R}^n \rightarrow \mathbb{R}^p$ which is a 1-cocycle for the action of $\pi_1(N)$ on $\text{Aff}(\mathbb{R}^n, \mathbb{R}^p)$ defined by $\gamma(C_3) = L_{2\gamma} \circ C_3 \circ (L_{1\gamma}, l_{1\gamma})^{-1}$. The bundle defined by the representation $h_\gamma^t(x, y) = (L_{1\gamma}(x) + l_{1\gamma}, L_{2\gamma}(y) + l_{2\gamma} + tC_3((L_{1\gamma}, l_{1\gamma})(x)) + l_{2\gamma})$, $0 \leq t \leq 1$ defines an homotopy between the bundle h_1^t defined by h^1 and the one h_0^t defined by h^0 . We deduce that h_0^t and h_1^t are isomorphic differentiable bundles. This implies that N_1 is diffeomorphic to the F -bundle defined by h^0 •

Let $(N, \nabla_F) = (F_0, \nabla_{F_0}), (F_1, \nabla_{F_1}), \dots, (F_n, \nabla_{F_n})$ be affine manifolds. We are going to define a gerbed tower which will allow us to study the classification of sequences $(N_n, \nabla_{N_n}) \rightarrow (N_{n-1}, \nabla_{N_{n-1}}) \rightarrow \dots \rightarrow (N, \nabla_N)$, where $(h_l : (N_l, \nabla_{N_l}) \rightarrow (N_{l-1}, \nabla_{N_{l-1}}))$ is an affinely locally trivial affine bundle).

Denote by T_{F_l} the group of translations of (F_l, ∇_{F_l}) . We suppose defined a flat $\text{Aff}(F_l, \nabla_{F_l})/T_{F_l}$ -bundle p_{F_l} over $(F_{l-1}, \nabla_{F_{l-1}})$. The bundle p_{F_l} induces a gerbe C_l defined over F_{l-1} (see proposition 4.3).

Definition 5.0.5.

We define $L_n \rightarrow L_{n-1} \dots \rightarrow L_1 \rightarrow L_0$ to be the sequence of 2-categories such that $L_1 \rightarrow L_0$ is C_1 , supposed defined L_p , an object of L_p is an affinely locally trivial (F_p, ∇_{F_p}) -bundle. A 1-arrow between a pair of objects e_p and e'_p of L_p is an affine map between their base space. A 2-arrow is an

isomorphism of affinely locally trivial bundle which cover a 1-arrow. We denote by $p_l : L_l \rightarrow L_{l-1}$ the projection.

An object of L_{p+1} is an affinely locally trivial $(F_{p+1}, \nabla_{F_{p+1}})$ -bundle e_{p+1} defined over an open subset U_p of an object e_p of L_p such that the restriction to each fiber of $U_p \rightarrow p_p(e_p)$ of the $Aff(F_{p+1}, \nabla_{F_{p+1}})/T_{F_{p+1}}$ -bundle induced, is the restriction of the $Aff(F_{p+1}, \nabla_{F_{p+1}})/T_{F_{p+1}}$ -bundle induced is $p_{F_{p+1}}$. A 1-arrow $h_{p+1}^1 : e_{p+1} \rightarrow e'_{p+1}$ between a pair of objects e_{p+1} and e'_{p+1} of L_{p+1} is a affine map between their respective base spaces e_p and e'_p induced by a 2-arrow of L_p . A 2-arrow between e_{p+1} and e'_{p+1} is an isomorphism of affinely locally trivial $(F_{p+1}, \nabla_{F_{p+1}})$ -bundles which covers a 1-arrow. We suppose that the 2-arrows depend only of the open subset $p_1 \dots p_{p+1}(e_{p+1})$ and $p_1 \dots p_{p+1}(e'_{p+1})$. Let h_p^2 be a 2-arrow of L_p , we define h_p^{2*} to be a 2-arrow of L_{p+1} which cover h_p^2 •

Proposition 5.0.6.

The sequence $L_n \rightarrow L_{n-1} \dots \rightarrow L_1 \rightarrow L_0$ that we have just defined is a gerbed tower.

Proof.

The fibered category $L_1 \rightarrow L_0$ is a gerbe as shows proposition 4.4. Let V be an open subset of N , and U an open subset of V , the restriction functor $r_{U,V} : L_{pU} \rightarrow L_{pV}$ is defined by the restriction of bundles.

Let U be an open subset of N , and e_l be an object of L_{lU} , the band of L_{l+1e_l} is the sheaf of sections of the $T_{F_{l+1}}$ -bundle p'_{e_l} defined on e_l induced by p_l , this sheaf does not depend of the objects chosen in the fibre L_{l+1e_l} , since we have supposed that the 2-arrows depend only of N .

Consider 2-morphisms $u_p : e_p \rightarrow e'_p$, and $u'_p : e'_p \rightarrow e''_p$ of L_p . We have defined in the paragraph above the proposition a morphism u_p^* . These morphisms satisfy $u'_p{}^* u_p^* = c(u_p, u'_p)(u'_p u_p)^*$, where $c(u_p, u'_p)$ is an automorphism of an object of the gerbe $L_{p+1e''_p}$ induced by the band •

7 Interpretation of the integral cohomology of a manifold.

Characteristic classes have been used by many mathematicians to study geometric objects. On this purpose, we have to give a geometric interpretation of the group $H^n(N, \mathbb{Z})$. This is what we propose to do in this part.

It is a well-known fact that the group $H^2(N, \mathbb{Z})$ is the set of equivalence classes of complex line bundles over N . Brylinski has defined an equivalence between the space of equivalence classes of complex line gerbes and $H^3(N, \mathbb{Z})$.

Consider $\mathcal{I}_N^*$ the sheaf of differentiable $\mathcal{I} - \{0\} = \mathcal{I}^*$ -functions defined on N . We say that a class $[c_n]$ of $H^n(N, \mathcal{I}_N^*)$ is geometric, if and only if there exists an $(n-1)$ -gerbed tower which classifying cocycle is c_n . A sufficient condition for a class c_n to be geometric is the following: there exists a classifying cocycle c_{n-1} of a commutative $(n-2)$ -gerbed tower $E(L_1, \dots, L_{n-2})$ which is an element of $H^{n-1}(N, L_{n-2})$, an exact sequence of sheaves $0 \rightarrow \mathcal{I}_N^* \rightarrow L \rightarrow L_{n-2} \rightarrow 0$ such that $[c_n]$ is the image of $[c_{n-1}]$ by the boundary map $\delta : H^{n-1}(N, L_{n-2}) \rightarrow H^n(N, \mathcal{I}^*)$.

We have the exact sequence

$$0 \rightarrow \mathbb{Z} \xrightarrow{i} \mathcal{I}_N \xrightarrow{exp} \mathcal{I}^*_N \rightarrow 0.$$

where i is the canonical injection, and $\exp$ the exponential map. It results from this sequence an isomorphism between $H^n(N, \mathcal{C}_N^*)$ and $H^{n+1}(N, \mathbb{Z})$. An element of $H^{n+1}(N, \mathbb{Z})$ will be said geometric if and only if it is the image of an element of $H^n(N, \mathcal{C}_N^*)$ which is the classifying cocycle of a $(n-1)$ -gerbed tower by the preceding isomorphism.

We have the following result:

Theorem 6.0.1.

Let N be a differentiable manifold, then each geometric class of $H^{n+2}(N, \mathbb{Z})$ is the classifying cocycle of a n -gerbed tower defined on N .

8 n -categories, and sheaves of n -categories.

In this part, we will define a notion of sheaf of n -categories over a topos N .

Definition 7.0.1.

A 0-pseudo-category is a set, a 1-pseudo-category C_1 , is a category.

Suppose defined the notion of n -pseudo-category.

An $(n+1)$ -pseudo-category C_{n+1} , is defined by, a class of objects $Ob(C_{n+1})$, for each objects x , and y , the n -pseudo-category of morphisms $Hom(x, y)$. For each objects u_1, u_2 and u_3 of C_{n+1} , there exists a composition n -functor:

$$Hom(u_2, u_3) \times Hom(u_1, u_2) \longrightarrow Hom(u_1, u_3)$$

We suppose the existence of an object 1_x of $Hom(x, x)$, such that for each arrow $h : x \rightarrow y$, $h \circ 1_x$ is isomorphic to h , and for each arrow $h' : y' \rightarrow x$, $1_x h'$ is isomorphic to h' .

An isomorphism between the objects x and y of an $n+1$ -pseudo-category is a map $f : x \rightarrow y$, such that there exists $h : y \rightarrow x$ such that hf is isomorphic to 1_x , and fh to 1_y .

A functor between two n -pseudo-categories C_n and C'_n , is defined as follows:

(i) A map $F : Ob(C_n) \rightarrow Ob(C'_n)$, and for each arrow $f : x \rightarrow y$, a morphism $F(f) : F(x) \rightarrow F(y)$ such that $F(f \circ f')$ is isomorphic to $F(f) \circ F(f')$.

(ii) A natural transformation between two functors F and F' , is defined by a family of maps $u_x : F(x) \rightarrow F'(x)$ such that for each map $f : x \rightarrow y$, $u_y F(f)$ is isomorphic to $F'(f) u_x$. At this stage, we do not precise the gluing datas.

Definition 7.0.2.

A 0-sheaf of sets defined on N , will be a sheaf of sets. Suppose defined the notion of sheaves of $n-1$ -pseudo-categories. A sheaf of n -pseudo-categories, will be defined by the following data: for each pair of objects U and V of N , and a map $h : U \rightarrow V$, a restriction functor $r^{C_n}_{U,V}(h) : C_n(V) \rightarrow C_n(U)$ such that for each triple of objects U_1, U_2 and U_3 , there exists an isomorphism $c(U_1, U_2, U_3)$ between $r^{C_n}_{U_1, U_2}(h) r^{C_n}_{U_2, U_3}(f)$ and $r^{C_n}_{U_1, U_3}(fh)$.

Gluing condition for objects:

Let $(U_i)_{i \in I}$ be an open cover of the object U of N , x_i an object of $C_n(U_i)$. If there exists, a family of maps $u_{jl} : r^{C_n}_{U_l \times_N U_j, U_l}(x_l) \rightarrow r^{C_n}_{U_l \times_N U_j, U_l}(x_j)$, a sheaf of $n-1$ -pseudo-categories

C_{n-1} defined on U , such that the restrictions map $r^{C_{n-1}}_{U_i \times_N U_j, U_i}$ are u_{ji} , then there exists an object x of $C_n(U)$ such that the restriction of x to U_i is x_i .

Gluing conditions for arrows:

For each x , and y in $C_n(U)$, the map defined on sub-objects of U by $V \rightarrow \text{Hom}(x|_V, y|_V)$ is a sheaf of $(n-1)$ -categories •

We denote $\mathcal{N}_n$ the pseudo-category whose objects are the elements of the set $\{1, \dots, n\}$, $\text{Hom}(j_1, j_2)$ has one element if j_1 inferior to j_2 , if not it is empty. We endow it with the topology such that the covering family of l are the integers inferior to l .

Definition 7.0.3.

An n -category, is a n -pseudo-category, such that for any objects $x_0, \dots, x_n, \dots$ of C , for each family of maps $u_{ij} : x_j \rightarrow x_i$ such that $u_{i_1 i_2} u_{i_2 i_3} = u_{i_1 i_3}$, the map defined on $\mathcal{N}$, $i \rightarrow \text{Hom}(x_i, x_0)$, is a sheaf of $n-1$ -pseudo-categories whose restrictions functors $u_{ij}^* : \text{Hom}(x_j, x_0) \rightarrow \text{Hom}(x_i, x_0)$ are defined by: $h \rightarrow hu_{ji}$ •

References

- [1] BENABOU, *Some remarks on 2-categorical algebra I*, Bul. Soc. Math. Belg., 41 (1989) pp. 127-194.
- [2] A. BOREL, *Linear algebraic groups*, W.A. Benjamin, Inc, New York-Amsterdam., (1969).
- [3] L. BREEN, *On the classification of 2-gerbes and 2-stacks*, Asterisque., 225 (1994).
- [4] G. E. BREDON, *Sheaf theory*, McGraw-HillBook Co., (1967).
- [5] J. L. BRYLINSKI, *J.L Loops spaces, Characteristic Classes and Geometric Quantization*, Progr. Math. 107, Birkhauser., (1993).
- [6] J. L. BRYLINSKI, AND D. A. MC LAUGHLIN, *The geometry of degree four characteristic classes and of line bundles on loop spaces I*, Duke Math. Journal., 75 (1994), pp. 603-637.
- [7] Y. CARRIERE, *Autour de la conjecture de L. Markus sur les variétés affines*, Invent. Math., 95 (1989), pp. 615-628.
- [8] P. DELIGNE, *Theorie de Hodge III*, Inst. Hautes Etudes Sci. Publ. Math., 44 (1974), pp. 5-77.
- [9] J. DUSKIN, *An outline of a theory of higher dimensional descent*, Bull. Soc. Math. Bel. Série A., 41 (1989), pp. 249-277.
- [10] D. FRIED, *Closed similarity affine manifolds*, Comment. Math. Helv., 55 (1980), pp. 576-582.
- [11] J. GIRAUD, *Cohomologie non abélienne*
- [12] D. FRIED AND W. GOLDMAN, *Three-dimensional affine crystallographic groups*, Advances in Math., 47 (1983), pp. 1-49.
- [13] D. FRIED, W. GOLDMAN, AND M. HIRSCH, *Affine manifolds with nilpotent holonomy*, Comment. Math. Helv., 56 (1981), pp. 487-523.

- [14] W. GOLDMAN, *Two examples of affine manifolds*, Pacific J. Math., 94 (1981), pp. 327-330.
- [15] W. GOLDMAN, *The symplectic nature of fundamental groups of surfaces*, Advances in Math., 54 (1984), pp. 200-225.
- [16] W. GOLDMAN, *Geometric structure on manifolds and varieties of representations*, 169-198, Contemp. Math., 74.
- [17] W. GOLDMAN AND M. HIRSCH, *The radiance obstruction and parallel forms on affine manifolds*, Trans. Amer. Math. Soc., 286 (1984), pp. 629-649.
- [18] W. GOLDMAN AND M. HIRSCH, *Affine manifolds and orbits of algebraic groups*, Trans. Amer. Math. Soc., 295 (1986), pp. 175-198.
- [19] R. GODEMENT, *Topologie algébrique et théorie des faisceaux* Herman., (1958).
- [20] J. L. KOSZUL, *Variétés localement plates et convexité*, Osaka J. Math., (1965), pp. 285-290.
- [21] A.-O. KUKU, *Ranks of K_n and G_n of orders and groups rings of finite groups over integer in number fields*, Journal of Pure and Applied Algebra., 138 (1999), 39-44.
- [22] S. MACLANE, *Homology*, Springer-Verlag., (1963).
- [23] G. MARGULIS, *Complete affine locally flat manifolds with a free fundamental group*, J. Soviet. Math., 134 (1987), pp. 129-134.
- [24] J. W. MILNOR, *On fundamental groups of complete affinely flat manifolds*, Advances in Math., 25 (1977), pp. 178-187.
- [25] M. Nguiffo Boyom, *Algebres a associateur symetrique et algebres reductives*, These Universite de Grenoble., (1968).
- [26] A. Tsemo, *Dynamique des variétés affines*, J. London Math. Soc., 63 (2001) pp. 469-487.
- [27] A. Tsemo, *Fibrés affines*, Michigan J. Math., 49 (2001) pp. 459-484.
- [28] A. Tsemo, *Composition series of affine manifolds and n -gerbes* Adv. Math. Research 3, Nova Sci. Publ., (2003), pp. 1-37.